



Roma, 29/01/2019

Ufficio: DOR/DMS
Protocollo: 201900001097/AG
Oggetto: **Articolo rivista scientifica analisi dati Progetto nazionale cefalee e farmacie di comunità.**
Circolare n. 11350
Sito Sì
4.1
Ifo sì

AI PRESIDENTI DEGLI ORDINI
DEI FARMACISTI
e p.c. AI COMPONENTI IL COMITATO
CENTRALE DELLA F.O.F.I.
LORO SEDI

**Progetto nazionale cefalee e farmacie di comunità:
pubblicato articolo scientifico che dimostra i risultati raggiungibili
con l'attività del farmacista attraverso il monitoraggio del corretto
uso della terapia nei pazienti cefalalgici.**

Si fa seguito e riferimento alle precedenti circolari federali aventi ad oggetto il Progetto nazionale cefalee e farmacia di comunità (cfr. circolari nn. 9563 del 30.10.2015 e 9594 del 20.11.2015), per fornire i seguenti aggiornamenti dell'iniziativa scientifica.

Come si ricorderà, il Progetto è coordinato dalla Prof.ssa Paola Brusa e coinvolge, oltre alla Federazione, il Dipartimento di Scienza e Tecnologia del Farmaco dell'Università di Torino, la Fondazione Italiana Cefalee (FICEF) e la Asl 3 di Torino. Le principali finalità dello studio scientifico erano la valorizzazione del ruolo del farmacista e lo sviluppo di nuove competenze professionali, da ottenere attraverso la dimostrazione dei risultati che la sua attività professionale e, in particolare, il monitoraggio del corretto uso della terapia, poteva apportare alla tutela della salute dei pazienti affetti da cefalalgia.

Le tappe principali del Progetto prevedevano:

- l'arruolamento di farmacisti di comunità sul territorio nazionale (si veda la citata circolare n. 9563, circa l'individuazione da parte degli Ordini di referenti operativi sul territorio);

- l'erogazione di un corso di Formazione a Distanza (si veda la suddetta circolare n. 9594, inerente l'attivazione della FAD che era gratuita per tutti i farmacisti iscritti all'Albo e consentiva l'acquisizione di 11 crediti ECM);
- la somministrazione di questionari in farmacia ai pazienti emicranici;
- la successiva analisi dei dati statistici ed epidemiologici emersi dai questionari.

La Federazione è lieta di informare che la Prof.ssa Brusa ha comunicato la recente pubblicazione di un secondo articolo scientifico, nel quale sono raccolti i dati emersi dallo studio ed i risultati raggiunti dal Progetto realizzato sotto l'egida federale (cfr. all. 1 e 2).

Tenuto conto della rilevanza scientifica dell'iniziativa e dei suoi riflessi in tema di promozione dell'attività professionale del farmacista, si invitato gli Ordini a diffondere presso gli iscritti i sopraindicati risultati del Progetto, attraverso i canali di comunicazione a propria disposizione.

Cordiali saluti.

IL SEGRETARIO
(Dr. Maurizio Pace)

IL PRESIDENTE
(On. Andrea Mandelli)

All. 2

RESEARCH ARTICLE

Self-medication for migraine: A nationwide cross-sectional study in Italy

Paola Brusa^{1,2*}, Gianni Allais^{3,4}, Cecilia Scarinzi⁵, Francesca Baratta¹, Marco Parente¹, Sara Rolando³, Roberto Gnani⁵, Teresa Spadea⁵, Giuseppe Costa⁵, Chiara Benedetto³, Massimo Mana⁶, Mario Giaccone^{2,7}, Andrea Mandelli⁷, Gian Camillo Manzoni⁴, Gennaro Bussone⁴

1 Department of Science and Technology of Drugs, University of Turin, Turin, Italy, **2** Order of Pharmacists of Turin, Turin, Italy, **3** Department of Surgical Sciences, Women's Headache Center, University of Turin, Turin, Italy, **4** FI.CEF Onlus, Italian Headache Foundation, Milan, Italy, **5** Epidemiology Unit, ASL TO3, Grugliasco (Turin), Italy, **6** ATF Informatics, Cuneo, Italy, **7** FOFI, Federation of the Orders of Italian Pharmacists, Rome, Italy

* paola.brusa@unito.it



Abstract

Headache disorders are considered the second leading cause of years lived with disability worldwide, and 90% of people have a headache episode at least once a year, thus representing a relevant public health priority. As the pharmacist is often the first and only point of reference for people complaining of headache, we carried out a survey in a nationwide sample of Italian pharmacies, in order to describe the distribution of migraine or non-migraine type headaches and medicines overuse among people entering pharmacies seeking for self-medication; and to evaluate the association, in particular of migraine, with socio-demographic and clinical characteristics, and with the pathway of care followed by the patients. A 14-item questionnaire, including socio-demographic and clinical factors, was administered by trained pharmacists to subjects who entered a pharmacy requesting self-medication for a headache attack. The ID Migraine™ Screener was used to classify headache sufferers in four classes. From June 2016 to January 2017, 4424 people have been interviewed. The prevalence of definite migraines was 40%, significantly higher among women and less educated people. About half of all headache sufferers and a third of migraineurs do not consider their condition as a disease and are not cared by any doctor. Among people seeking self-medication in pharmacies for acute headache attacks, the rate of definite or probable migraine is high, and a large percentage of them is not correctly diagnosed and treated. The pharmacy can be a valuable observatory for the study of headaches, and the first important step to improve the quality of care delivered to these patients.

OPEN ACCESS

Citation: Brusa P, Allais G, Scarinzi C, Baratta F, Parente M, Rolando S, et al. (2019) Self-medication for migraine: A nationwide cross-sectional study in Italy. PLoS ONE 14(1): e0211191. <https://doi.org/10.1371/journal.pone.0211191>

Editor: Md. Asiful Islam, Universiti Sains Malaysia, MALAYSIA

Received: October 24, 2018

Accepted: January 8, 2019

Published: January 23, 2019

Copyright: © 2019 Brusa et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the manuscript.

Funding: The authors declare that this study was funded by FOFI, Federation of the Orders of Italian Pharmacists, Rome, Italy. The source of the funding had no role in the design of the study or the analysis and interpretation of the data. The role of the Federation was solely to verify that the study had been conducted in accordance with the agreed methodology.

Introduction

Headache disorders are considered by The Global Burden of Disease study as the second leading cause of years lived with disability worldwide [1]. The World Health Organization declared that headache disorders are ubiquitous, prevalent and disabling but despite this

Competing interests: The authors have declared that no competing interests exist.

Abbreviations: CM, chronic migraine; EM, episodic migraine; ID-MS, ID Migraine™ Screener; IHS, International Headache Society; MOH, medication overuse headache; MOs, medication overusers; NSAID, nonsteroidal anti-inflammatory drug; OTC, over the counter; TM, transformed migraine.

they are under-recognized, under-diagnosed and under-treated all over the world [2]. Ninety percent of people have a headache episode at least once a year and 40% are serious or disabling headaches. Headaches cause a loss of productivity at work, a reduction in social interactions and a deterioration in the quality of life [2]. Globally, it has a prevalence of 47% and is the most common neurological syndrome for which people turn to their family doctor while only 16% consult a neurologist [2]. Recent epidemiological studies have shown that the phenomenon of migraine may be compared to an iceberg: the top represents correctly diagnosed cases while the submerged part is made up of those subjects that have never consulted a doctor or that, having done so, have not received a correct diagnosis [3, 4]. Furthermore, in Italy, many migraineurs make a wrong self-diagnosis assuming cervical spine pathology as the cause [5]. This explains why the majority of migraineurs make use of self-medication, taking over the counter (OTC) analgesics to treat their headache [6,7]. The lack of a proper treatment for headache, however, can lead to an overuse of acute pain-relief medicine and to the medication overuse headache (MOH), an even more disabling chronic disorder [8,9].

In Europe, the total annual cost of headache amongst adults aged 18–65 years was calculated, in a study of 2012, at €173 billion [10]. In Italy, among the neurological disorders, migraine ranked first for direct and indirect costs with a total expenditure amounting to € 3.5 billion per year [11]. An Italian study conducted on 135 patients with MOH, demonstrated that the direct costs, calculated by medications for acute treatment and prophylaxis, diagnostic procedures, visits, complementary treatments, informal care, were around 415 €/month per person; indirect costs were 530 €/month, and were mostly due to presenteeism (350 €, 66.3%), i.e. days worked with headache, rather than to absenteeism, lost workdays (160 €, 33.7%) [12]. Thus, headaches in general and migraines in particular represent not only an enormous limitation for the affected person, but a problem for the entire community [13].

The pharmacist is often the first and only point of reference for the patient complaining of headache, and participates in the management of this widespread disorder. He/she should be able to understand the problem raised by the patient, recommend the most appropriate symptomatic treatment and, if necessary, refer the patient to a general practitioner or a specialist. With proper training, the community pharmacist could be responsible for identifying patients suffering from suspected migraine symptoms that have not been diagnosed, owing to poor awareness about the problem and lack of specialist advice [14]. Community pharmacies are also the ideal site where the risks associated with inappropriate self-medication could be prevented. Community pharmacists are strategically positioned as they have a clear overview of the medications taken by a patient, whether prescribed by a physician or purchased directly [15].

Considering the potential role of community pharmacists, and his/her strategical role as first contact with headache sufferers, a preliminary study was carried out in Piedmont in 2014, with the objective of analyzing the patterns of use of drugs in subjects who entered a pharmacy requesting OTC medications in order to treat a headache attack [16]. Thanks to the findings of that study, both from an epidemiological perspective and a public health policy point of view [16, 17], the decision was taken to expand the scale of the model to a national level. The detailed protocol of the national survey has already been reported [18]. The objectives of the present paper are to describe the distribution of migraine or non-migraine type headaches and medicines overuse among people entering pharmacies seeking for self-medication; and to evaluate the association, in particular of migraine, with socio-demographic and clinical characteristics, and with the pathway of care followed by the patients.

Materials and methods

The study was designed as a cross-sectional survey based on face-to-face interviews using questionnaires drawn up by experts and based on the scientific literature. Briefly, the questionnaire, reported in Fig 1, covered the following points: the socio-economic situation of the recruited

- 1) Sex M F
- 2) Age [actual age] _____
- 3) Height [in cm] and weight [in kg] _____
- 4) Education [the highest qualification attained]
 - University degree or higher _____
 - 4-5-year High School Diploma valid for third-level education _____
 - 2-3-year High School Diploma not valid for third-level education _____
 - Junior High School Diploma (or apprenticeship/job) _____
 - Primary school exam _____
 - No qualification _____
- 5) Occupation

If employed:

 - Manual labourer _____
 - Employee non-manual (conceptual/intellectual) _____
 - Self-employed:
 - Businessperson _____
 - Professional _____
 - Self-employed worker _____

Unemployed _____

 - Homemaker _____
 - Student _____
 - Retired _____
 - Other (military service, invalid etc.) _____
- 6) How many members are there in your family? Number of people in your household (including patient) _____
- 7) ID Migraine screener test
 - Did you have the following with your headaches:
 - You felt nauseated or sick to your stomach? YES NO
 - Light bothered you (a lot more than when you didn't have headaches)? YES NO
 - Your headaches limited your ability to work, study, or do what you needed to do for at least 1 day? YES NO

If the patient answers YES to at least three questions, there is a 95% probability that he/she suffers from migraines
- 8) How long have you been having headaches?
 - <1 year; 1-4 years;
 - 5-9 years; ≥10 years.
- 9) How many days have you had a headache in the last 3 months? n° _____
- 10) How many days each month do you take pain medication for a headache?
 - up to 3; from 4 to 10; more than 10.
- 11) Does the headache go away after you have taken pain medication?
 - Yes; Often; Rarely; No.
- 12) Which medicines do you usually take when you have an acute attack? [more than one answer is permitted]
 - nonsteroidal anti-inflammatory drugs (NSAIDS);
 - ergot-derivatives;
 - tryptans;
 - combination drugs;
 - other analgesics
- 13) Are you being treated by a medical professional? [more than one answer is allowed]
 - nobody;
 - my family doctor;
 - a specialist;
 - a headache centre
- 14) Do you regard your headache as an illness? YES NO.

Fig 1. Questionnaire.

<https://doi.org/10.1371/journal.pone.0211191.g001>

subjects; the ID Migraine screener test (ID-MS) [19,20]; the frequency of attacks, the drugs usually taken and the healthcare professional, if any, responsible for the management of the subject's headache.

Given the nature of the centres responsible for recruiting the subjects (community pharmacies), a single national online training course was employed to prepare the pharmacists in order to ensure that the data were collected as homogeneously as possible. At the end of the training course, each pharmacist had to pass a final test before being enrolled in the project. Pharmacists were selected on a voluntary basis, and did not receive any payment for their participation in the survey.

The questionnaire, was administered to subjects who entered a pharmacy requesting self-medication for a headache attack, from June 2016 to January 2017. The subjects were recruited at a maximum rate of five patients per pharmacy per month. [18]

Age at enrolment was categorized in three levels: less than 30 years, 30–54 years and more than 54 years. Educational level was measured by means of the maximum attained classification and classified according to the International standard classification [21] in three levels: low, including people up to the completion of lower secondary school (ISCED 0–2); medium, including higher secondary school diplomas (ISCED 3); and high, with any university degree (ISCED 5–6).

The clinical section included information about symptoms, occurrence and severity of headaches, together with the items of the ID-MS, resulting in a four-level variable that classifies patients as suffering from definite migraine, probable migraine, unlikely migraine, or other headaches [19, 20]. The variable “Number of days with headache in the last three months” was categorized in two ways:

1. 0–44 days and ≥ 45 days in accordance with the criteria laid down by the International Headache Society (IHS) classification [22] to distinguish episodic migraine (EM) (<15 days per month) from chronic migraine (CM) (≥ 15 days per month);
2. <11 days, 11–29 days, 30–60 days e > 60 days in line with the time limits recently recommended by Manzoni et al [23] in order to distinguish respectively: infrequent EM (≤ 3 days per month), frequent EM (> 3 but < 10 days per month), very frequent migraine—or CM—(≥ 10 but ≤ 20 days per month) and transformed migraine (TM) (> 20 days per month).

People who declared to take pain medications for more than 10 days a month were classified as medicines overusers (MOs).

Finally, as for pathway of care, patients were asked which medicines were usually taken in case of an acute attack and whether they were monitored by a professional for the treatment of headache. More details on the protocol of the survey are reported elsewhere [18].

To describe the association of age, pathway of care and clinical information with the type of headache, we computed conditional relative frequencies and related chi-square p-values, stratified by gender. The Monte-Carlo exact test was applied if the cells with expected conjoint frequencies lower than 5 were more than 20%. Statistical analyses were carried out using the SAS software (version 9.2).

Patient involvement

The study was designed in order to describe the distribution of migraine or non-migraine type headaches and medicines overuse among people seeking for self-medication, and to evaluate its association with individual characteristics and with the pathway of care followed by the patients. The study was approved by the national association of cephalalgic patients. The study was also approved and then financed by the Federation of the Orders of Italian Pharmacists

(FOFI). Approval by an ethics committee was not required because the subjects participated in the study on a voluntary basis and they were informed on the characteristics and the purpose of the study. The questionnaire was anonymous, personal data were not collected and there is no way to trace back the answers to a specific responder.

Results

The development of the study at a national level expanded the number of participating pharmacists to 610, operating in 514 community pharmacies situated in 19 out of the 20 Italian regions. Out of 610 trained pharmacists, 445 collected at least one questionnaire and participated in this survey; overall, 4424 questionnaires had been correctly completed.

The prevalence of migraine and headache according to the ID-MS classification, by socio-demographic characteristics is summarized in Table 1. The study population included mostly women (74%), highly educated people (only 25% in the lowest level), residing in the North of Italy (84%), and with a mean age of 45 years. Among people asking for self-medication because of headache, 40% has a definite migraine and 31% a probable migraine, while the remaining 30% is unlikely to have a migraine or has other forms of headaches.

The ID-MS shows a strong association with gender: almost half of the women (45%) were classified as definite migraine vs. only a quarter of men (26%), while the percentage of men with other types of headaches was twice that of women (14% vs. 7%). The overall ratio of females to males equals 2.8:1, but it is 4.9:1 among subjects with definite migraine and 1.4:1 for non-migraine sufferers.

Within the sample, the rate of definite migraines was significantly higher among the two lower educational levels than among people with university qualifications (44% and 42% vs. 35%); there were slightly more other headaches in people living in the North compared to the rest of Italy (9% vs. 6%). Mean age was homogeneously distributed across the classes of migraine according to the ID-MS classification.

Table 1. Prevalence (%) of migraine according to the ID-MS classification by socio-demographic characteristics.

Socio-demographic characteristics	migraine classes according to the ID-MS classification				Total		p-value §
	other headaches (N = 389)	unlikely migraine (N = 898)	probable migraine (N = 1372)	definite migraine (N = 1765)	N	%	
	%	%	%	%			
Gender							
Female	7.0	18.1	29.9	45.0	3265	73.8	<0.0001*
Male	13.7	26.4	34.3	25.6	1159	26.2	
Educational level							
Low	8.2	20.1	28.3	43.4	1112	25.1	0,0001*
Medium	8.9	18.7	30.7	41.7	1900	42.9	
High	9.1	22.7	33.5	34.7	1412	31.9	
Residence area							
North	9.3	19.6	31.3	39.8	3726	84.2	0,004*
Center, South and Isles	6.2	24.1	29.4	40.4	698	15.8	
Age in years						mean value	
mean	46.0	45.5	44.6	45.1		45.1	0.21
Total (%)	8.8	20.3	31.0	39.9	4424		

§ p-values referred to the chi-square or Monte-Carlo exact tests on the distribution of the socio-demographic characteristics across all ID-MS classes

* Statistically significant

When looking at men and women separately (Table 2), a chronic headache (at least 45 headache days over the previous three months) is reported in 5.4% of men ($n = 63$) and in 7.9% of women ($n = 257$) and MOs in 14.9% ($n = 173$) and 18.8% ($n = 613$), respectively. Less than 10% (130 men and 189 women) have had this condition for less than one year, while about 60% have been affected for at least 10 years (563 men and 2039 women).

Almost all the covariates are significantly associated with the type of headache, except for age and some types of drugs (as ergot-derivatives, combination drugs and other analgesics). In particular, a higher duration and frequency of headache (higher number of years and of days in each month) is associated with higher prevalence of “definite migraine”, as well as a higher number of drugs. These associations are particularly evident among women. Also MOs were significantly overrepresented among subjects with definite or probable migraine (76% among men and 88% among women); only 6.4% of men MOs and 2.4% of women suffered from non-migraine headaches.

As for therapies, triptans and nonsteroidal anti-inflammatory drug (NSAID) users are mostly affected by more severe migraines (81% and 70%, respectively among men; 91% and 84% among women), but only 38% of subjects with definite migraine take triptans in order to block the attack. On the other hand, men use all the other types of drugs independently of the migraine class, while women with other forms of headaches tend to use only other analgesics. Headaches classified as probable or definite migraines are more often not resolved by pain medication, other types of headaches are usually blocked with symptomatic medications.

Finally, about 50% of all patients and 30% of those who suffer from definite migraines do not believe that headache is a disease and do not require medical care, especially among men. Indeed, 49% of men and 63% of women who are not cared by any doctor are likely to be serious migraineurs. Of those who do consult a doctor, half goes to the general practitioner and the other half is in care with a specialist or at a headache centre. For both genders, the higher the ID-MS class, the higher the percentage of patients who believe that migraine is a disease and who are in care by either a physician or a Headache Center.

Discussion

In accordance with other studies [13] we report that women are more likely than men to suffer from headaches, particularly from migraines. In both genders, there appears to be a trend towards mainly non-occasional headaches and migraines, existing for a long time and accompanied by an important prevalence of chronic headache. Since the epidemiological data reported in the literature indicate a prevalence of chronic migraine in the general population of only 2%, must be true that the migraineurs entering a pharmacy and enrolled in this survey have a relatively severe form of migraine.

Despite the specificity and usefulness of triptans, which have been available on the market for over 25 years, we found that only one third of severe migraineurs use this class of medication. This is in line with data from Fisher *et al.*, where it has been reported that almost three quarters of patients admitted for a preliminary observation at the Headache Outpatient Clinic had never taken triptans before [24].

It has long been known that the overuse of symptomatic medications can worsen migraine, until it loses the typical episodic temporal pattern and transform it into a chronic headache [25]. In this context we can speak of MOs. In our study, approximately one quarter of subjects with definite migraine declared that they took symptomatic medications at a frequency deemed to be medication overuse by the criteria in the IHS classification [26]. If we consider the worsening of the migraine due to medication overuse and the difficulty in detecting migraineurs who often self-medicate without consulting a doctor, this high percentage of

Table 2. Prevalence (%) of migraine according to the ID-MS classification by gender and clinical characteristics.

Clinical characteristics	MALE						FEMALE					
	N	migraine classes				p-value §	N	migraine classes				p-value §
		other headaches N = 159	unlikely migraine N = 306	probable migraine N = 397	definite migraine N = 297			other headaches N = 230	unlikely migraine N = 592	probable migraine N = 975	definite migraine N = 1468	
	%	%	%	%		%	%	%	%			
Age												
≤29	138	11,6	27,5	34,1	26,8	0,67	447	6,7	18,3	33,3	41,6	0,16
30–54	724	13,8	25,7	33,4	27,1		2092	6,5	18,0	29,3	46,2	
≥55	297	14,5	27,6	36,4	21,5		726	9,0	18,3	29,3	43,4	
Number of years with headache												
<1	130	36,9	31,5	22,3	9,2	<0.0001*	189	20,6	37,0	26,5	15,9	<0.0001*
1–4	268	14,6	30,6	34,3	20,5		537	10,2	25,0	32,6	32,2	
5–9	198	11,1	22,7	38,9	27,3		500	6,8	16,8	33,8	42,6	
≥10	563	8,9	24,5	35,3	31,3		2039	5,0	14,9	28,5	51,6	
Number of days with headache in the last three months												
<45	1096	14,2	26,7	34,3	24,7	0,01*	3008	7,3	18,6	30,3	43,9	0,0002*
≥45	63	4,8	20,6	33,3	41,3		257	3,9	13,2	25,3	57,6	
Number of days with headache in the last three months												
<11	871	15,8	28,6	33,6	21,9	<0.0001*	2153	9,0	20,9	30,5	39,6	<0.0001*
11–29	177	6,8	22,0	35,6	35,6		638	2,8	12,9	29,2	55,2	
30–60	74	9,5	13,5	40,5	36,5		333	4,5	11,1	30,0	54,4	
>60	37	5,4	21,6	29,7	43,2		141	2,8	15,6	23,4	58,2	
Number of types of medicines												
1	935	15,1	28,2	33,9	22,8	<0.0001*	2531	8,0	20,3	30,9	40,8	<0.0001*
2	197	8,6	18,8	35,5	37,1		620	4,5	11,1	27,9	56,5	
≥3	27	3,7	18,6	37,0	40,7		114	0,0	7,9	17,5	74,6	
Medication Overuse Headache												
No	986	15,0	27,9	34,2	22,9	<0.0001*	2652	8,1	20,0	30,8	41,0	<0.0001*
Yes	173	6,4	17,9	34,7	41,0		613	2,4	10,0	25,6	62,0	
Type of drug												
NSAIDs	272	9,9	20,6	31,6	37,9	<0.0001*	964	3,4	12,6	25,1	58,9	<0.0001*
Ergot-derivatives	28	7,1	28,6	28,6	35,7		123	1,6	8,1	23,6	66,7	
Triptans	220	4,1	15,0	33,6	47,3	<0.0001*	855	1,5	7,0	24,9	66,5	<0.0001*
Combination drugs	82	9,8	20,7	36,6	32,9		284	2,5	14,1	20,8	62,7	
Other analgesics	197	13,7	22,8	34,0	29,4	0,48	564	6,9	17,6	27,3	48,2	0,35
Pain medication solved the problem												
Yes	613	18,9	32,8	32,0	16,3	<0.0001*	1384	11,8	25,1	32,6	30,6	<0.0001*
Often	360	10,0	19,7	39,4	30,8		1055	3,7	16,0	29,8	50,5	
Rarely	128	5,5	16,4	29,7	48,4		538	3,7	8,0	25,8	62,5	
No	58	0,0	22,4	36,2	41,4		288	2,8	11,5	24,7	61,1	

(Continued)

Table 2. (Continued)

Clinical characteristics	MALE						FEMALE					
	migraine classes						migraine classes					
	other headaches N = 159	unlikely migraine N = 306	probable migraine N = 397	definite migraine N = 297			other headaches N = 230	unlikely migraine N = 592	probable migraine N = 975	definite migraine N = 1468		
The headache is a disease												
No	677	20,4	33,1	34,4	12,1	<0.0001*	1545	12,4	26,9	33,7	27,0	<0.0001*
Yes	482	4,4	17,0	34,0	44,6		1720	2,2	10,3	26,4	61,1	
Who takes care of the patient												
Nobody	676	17,9	32,7	32,5	16,9	<0.0001*	1547	10,9	26,2	32,6	30,3	<0.0001*
General practitioner	318	8,2	18,6	39,3	34,0	<0.0001*	1030	4,2	12,1	27,6	56,1	<0.0001*
Specialist	135	8,1	12,6	35,6	43,7	<0.0001*	533	2,8	10,1	25,9	61,2	<0.0001*
Headache Center	82	3,7	20,7	32,9	42,7	0,0005*	365	2,7	5,5	24,1	67,7	<0.0001*

§ p-values referred to the chi-square or Monte-Carlo exact tests on the distribution of the clinical characteristics across all ID-MS classes

* Statistically significant

<https://doi.org/10.1371/journal.pone.0211191.t002>

migraineurs MO entering a pharmacy is clear evidence of the fundamental importance of this site as the first point of contact for medicines overuse management.

About half the subjects included in our study believe that their headache is not a disease and manage their pain by themselves without consulting a medical professional. The migraineurs have a greater awareness of this condition as a disease, and are more likely to be treated by medical professionals, although a third of subjects with definite migraine does not receive any medical consultation, thus giving wide space to interventions aimed to improve the quality and appropriateness of care delivered to these patients. The fact that this condition mainly affects middle-aged subjects very busy from the working point of view and generally in good health except for migraine problems, reduces the chance of consulting a doctor. An interesting study conducted in the US in 1998 [27] indicated that a mere 50% of migraineurs in the general population were in care with a doctor. They found, moreover, that even amongst those who consulted a doctor, the majority either had been misdiagnosed, or had been diagnosed correctly but were being treated with an inappropriate therapy or, even having been diagnosed correctly and receiving a correct therapy, had not been going back for regular follow-up examinations. The study arrived at the bitter conclusion that a tiny percentage of migraineurs (6%) were managed appropriately. The higher percentage of migraineurs consulting a doctor found by our study is probably due to the fact that not the general population was analysed but subjects were entered into a pharmacy looking for pain relief and, therefore, affected by a more severe form of migraine.

To sum up, the results of our investigation highlight that females make up the great majority of subjects entering a pharmacy seeking pain relief; approximately 40% of all cases were subjects with definite migraine and for many of them the migraine symptoms had been present for many years and with frequent attacks, which did not always respond to symptomatic therapies. About half of headache sufferers and about a third of migraineurs do not consider their condition as a disease and tend to self-medicate, which leads to inappropriate treatment of the condition.

A limit of our study is that the participating pharmacists were enrolled into the study on a volunteer basis, and are not a representative sample of the Italian population of pharmacists, as the rate of participation was much higher in Northern regions of Italy compared to Central or Southern. Consequently, patients interviewed cannot be considered as representative of the Italian population seeking for self medication in a pharmacy. Unfortunately, we cannot estimate the magnitude of this source of bias, as there are no available data to compare the characteristics of population entering in pharmacy with the general population. On the other hand, the fact that all interviewing pharmacists were volunteer and underwent a formal training in interviewing headache suffering customers should ensure a high level of quality of data collected.

An original contribution of our work is that we showed the strength of the pharmacies as epidemiological sentinels, given their capillary network on the field, and their contact with a wide range of users. This allows any pharmacy-based study to develop a snapshot close to the real-life situation and far more accurate than that obtained from data based in specialist centres, enabling a choice of clinical strategies that are coherent with the real needs of the local population.

Conclusion

The pharmacy can be not only a valuable observatory for the study of headaches, but can also represent, through the counselling that a formed pharmacist might give, the first important step in seeking to build a good management relationship for subjects with headache, especially so for undiagnosed migraineurs; their referral in the first instance to the general practitioner and then to specialists and headache centres, may prevent the onset of chronic migraine and medication overuse headache.

Acknowledgments

The authors would like to thank the Federation of the Orders of Italian Pharmacists (FOFI) for its contribution both to the extension of the project to the national level and financial support and the Italian Headache Foundation (FICEF Onlus) for its contribution in drawing up the project and the organisation of the online training course for pharmacists that participated in the study.

The authors also wish to thank Eng. Ezio Festa (ATF Informatics, Cuneo, Italy) for developing the data gathering software and database management.

We acknowledge the invaluable work of all the pharmacists who participated in carrying out the survey.

The authors also wish to thank Dr. Tom O Byrne for the linguistic revision of the text.

Author Contributions

Conceptualization: Paola Brusa, Gianni Allais.

Data curation: Cecilia Scarinzi, Sara Rolando, Roberto Gnavi, Teresa Spadea, Giuseppe Costa, Chiara Benedetto, Gian Camillo Manzoni, Gennaro Bussone.

Formal analysis: Cecilia Scarinzi, Sara Rolando, Roberto Gnavi, Teresa Spadea, Giuseppe Costa, Chiara Benedetto.

Funding acquisition: Paola Brusa.

Investigation: Francesca Baratta, Marco Parente, Sara Rolando, Chiara Benedetto.

Methodology: Gianni Allais, Giuseppe Costa.

Project administration: Paola Brusa.

Software: Massimo Mana.

Supervision: Paola Brusa, Mario Giaccone, Andrea Mandelli.

Writing – original draft: Paola Brusa, Gianni Allais, Francesca Baratta, Roberto Gnani, Teresa Spadea.

Writing – review & editing: Gian Camillo Manzoni, Gennaro Bussone.

References

1. Saylor D, Steiner TJ. The Global Burden of Headache. *Semin Neurol*. 2018 Apr; 38(2):182–190. <https://doi.org/10.1055/s-0038-1646946> PMID: 29791944
2. World Health Organization, Lifting the Burden (2011) Atlas of headache disorders and resources in the world. 2011. Geneva: WHO.
3. Dodick DW, Tepper SJ, Lipton RB, Buse DC, Stewart WF, Bayliss M et al. Improving Medical Communication in Migraine Management: A Modified Delphi Study to Develop a Digital Migraine Tracker. *Headache*. 2018 Oct; 58(9):1358–1372. <https://doi.org/10.1111/head.13426> PMID: 30362524
4. Miller S, Matharu MS. Migraine is underdiagnosed and undertreated. *Practitioner*. 2014 Sep; 258(1774):19–24. PMID: 25588281
5. Viana M, Sances G, Terrazzino S, Sprenger T, Nappi G, Tassorelli C. When cervical pain is actually migraine: An observational study in 207 patients. *Cephalalgia*. 2018 Feb; 38(2):383–388. <https://doi.org/10.1177/0333102416683917> PMID: 27927894
6. Katsarava Z, Mania M, Lampl C, Herberhold J, Steiner TJ. Poor medical care for people with migraine in Europe—evidence from the Eurolight study. *J Headache Pain*. 2018; 19(1):10–13 <https://doi.org/10.1186/s10194-018-0839-1> PMID: 29392600
7. Cevoli S, D'Amico D, Martelletti P, Valguarnera F, Del Bene E, De Simone R et al. Underdiagnosis and undertreatment of migraine 85 in Italy: a survey of patients attending for the first time 10 headache centres. *Cephalalgia*. 2009 Dec; 29(12):1285–93. <https://doi.org/10.1111/j.1468-2982.2009.01874.x> PMID: 19438916
8. Vandebussche N, Laterza D, Lisicki M, Lloyd J, Lupi C, Tischler H et al. Medication-overuse headache: a widely recognized entity amidst ongoing debate. *J Headache Pain*. 2018 Jul 13; 19(1):50. <https://doi.org/10.1186/s10194-018-0875-x> PMID: 30003412
9. Martelletti P. The journey from genetic predisposition to medication overuse headache to its acquisition as sequela of chronic migraine. *J Headache Pain* 2018 Jan 10; 19(1):2. <https://doi.org/10.1186/s10194-017-0830-2> PMID: 29322261
10. Linde M, Gustavsson A, Stovner LJ, Steiner TJ, Barré J, Katsarava Z et al. The cost of headache disorders in Europe: the Eurolight project. *Eur J Neurol*. 2012 May; 19(5):703–11. <https://doi.org/10.1111/j.1468-1331.2011.03612.x> PMID: 22136117
11. Pugliatti M, Sobocki P, Beghi E, Pini S, Cassano GB, Altamura AC et al. Cost of Disorders of the Brain in Europe Study Group. Cost of disorders of the brain in Italy. *Neurol Sci*. 2008 Apr; 29(2):99–107. <https://doi.org/10.1007/s10072-008-0868-7> PMID: 18483707
12. D'Amico D, Grazi L, Curone M, Leonardi M, Raggi A. Cost of medication overuse headache in Italian patients at the time-point of withdrawal: a retrospective study based on real data. *Neurol Sci*. 2017 May; 38(Suppl 1):3–6. <https://doi.org/10.1007/s10072-017-2891-z> PMID: 28527071
13. Manzoni GC, Stovner LJ. Epidemiology of headache. *Handb Clin Neurol* 2011; 97: 3–22.
14. Giaccone M, Baratta F, Allais G, Brusa P. Prevention, education and information: the role of the community pharmacist in the management of headaches. *Neurol Sci* 2014; 35:S1–S4.
15. Horvat N, Koder M, Kos M. Using the simulated patient methodology to assess paracetamol-related counseling for headache. *PLoS One* 2012; 7(12): e52510. <https://doi.org/10.1371/journal.pone.0052510> PMID: 23300691
16. Brusa P, Allais G, Bussone G, Rolando S, Giaccone M, Aguggia M et al. Migraine attacks in the pharmacy: a survey in Piedmont, Italy. *Neurol Sci* 2014; 35: S5–S9
17. Brusa P, Allais G, Rolando S, Baratta F, Giaccone M, Bussone G et al. Migraine attacks in the pharmacy: a gender subanalysis on treatment preferences. *Neurol Sci* 2015; 36: S93–S95

18. Brusa P, Parente M, Allais G, Rolando S, Costa G, Gnani R et al. Community pharmacies as epidemiological sentinels of headache: first experience in Italy. *Neurol Sci* 2017; 38(Suppl 1): 15–20. <https://doi.org/10.1007/s10072-017-2908-7> PMID: [28527066](https://pubmed.ncbi.nlm.nih.gov/28527066/)
19. Lipton RB, Dodick D, Sadovsky R, Kolodner K, Endicott J, Hettiarachchi J et al. A self-administered screener for migraine in primary care. The ID Migraine™ validation study. *Neurology* 2003; 61: 375–382 PMID: [12913201](https://pubmed.ncbi.nlm.nih.gov/12913201/)
20. Rapoport AM, Bigal ME. ID-migraine. *Neurol Sci* 2004; 25:S258–S260. <https://doi.org/10.1007/s10072-004-0301-9> PMID: [15549552](https://pubmed.ncbi.nlm.nih.gov/15549552/)
21. UNESCO Institute for Statistics. International Standard Classification of Education ISCED 2011. Montreal, Quebec, 2012. <http://uis.unesco.org/sites/default/files/documents/international-standard-classification-of-education-isced-2011-en.pdf>
22. Headache Classification Committee of the International Headache Society: The international classification of headache disorders, 3rd edition (beta version). *Cephalalgia* 2013, 33:629–808. <https://doi.org/10.1177/0333102413485658> PMID: [23771276](https://pubmed.ncbi.nlm.nih.gov/23771276/)
23. Manzoni GC, Bonavita V, Bussone G, Cortelli P, Narbone MC, Cevoli S et al. On behalf of ANIRCEF (Associazione Neurologica Italiana Ricerca Cefalee): Chronic migraine classification: current knowledge and future perspectives. *J Headache Pain* 2011, 12:585–592. <https://doi.org/10.1007/s10194-011-0393-6> PMID: [22028184](https://pubmed.ncbi.nlm.nih.gov/22028184/)
24. Fischer M, Frank F, Wille G, Klien S, Lackner P, Broessner G. Triptans for acute migraine headache: current experience with triptan use and prescription habits in a tertiary care headache outpatient clinic: an observational study. *Headache* 2016; 56:952–960. <https://doi.org/10.1111/head.12820> PMID: [27091593](https://pubmed.ncbi.nlm.nih.gov/27091593/)
25. Mathew NT, Stubits E, Nigam MP. Transformation of episodic migraine into daily headache: analysis of factors. *Headache* 1982; 22:66–68. PMID: [7085263](https://pubmed.ncbi.nlm.nih.gov/7085263/)
26. Headache Classification Committee of the International Headache Society (IHS). The International classification of headache disorders, 3rd edition. *Cephalalgia*. 2018 Jan; 38(1):1–211. <https://doi.org/10.1177/0333102417738202> PMID: [29368949](https://pubmed.ncbi.nlm.nih.gov/29368949/)
27. Lipton RB, Stewart WF, Simon D. Medical consultation for migraine: results from the American Migraine Study. *Headache* 1998; 38:87–96. PMID: [9529763](https://pubmed.ncbi.nlm.nih.gov/9529763/)

Community pharmacies as epidemiological sentinels of headache: first experience in Italy

Paola Brusa^{*,†}, Marco Parente^{*}, Giovanni Allais^{‡§}, Sara Rolando[^], Giuseppe Costa⁺, Roberto Gravi[†], Teresa Spaccat[†], Mario Giaccione[¶], Andrea Mandelli[¶], Massimo Mana^{¶¶}, Francesca Baratta^{*}, Chiara Benocetto[^], Gianro Bussone[§]

^{*} Department of Scienza e Tecnologia del Farmaco, University of Turin, Italy

[†] Order of Pharmacists of Turin, Italy

[^] Department of Surgical Sciences, Women's Headache Center, University of Turin, Italy

[§] Ficef Onlus, Milan, Italy

[†] Epidemiology Unit, ASL TO3, Grugliasco, Italy

[¶] FOF, Federation of the Orders of Italian Pharmacists, Rome, Italy

^{¶¶} ATF informatics, Cuneo, Italy

^{††} Federfarma Piemonte, Turin, Italy

Keywords:

Community pharmacy, medication overuse headache, migraine.

Abbreviations:

MOH medication overuse headache, OTC over-the-counter.

Conflicts of interest:

All the authors certify that there is no actual or potential conflict of interest in relation to this article.

Acknowledgements:

The authors would like to thank Federation of the Orders of Italian Pharmacists (FOFI) that funded the national project.

Summary

Migraine is a disabling neurovascular syndrome which affects 12-15% of the global population and it represents the third cause in years lived with disability in both males and females aged 15-49 years.

Among migraineurs, the symptomatic drug abuse may be a risk factor in the development of medication overuse headache (MOH).

Detecting cases of MOH is not straightforward; community pharmacists may therefore be in a strategic position to identify individuals who self-medicate, particularly with respect to prevent the development of MOH.

In 2014, our group published the results of a survey conducted in Piedmont, Italy, on the patterns of use and dispensing of drugs in patients requesting assistance from pharmacists for relief of a migraine attack.

We decided, now, to expand the scope of the model to a national level. The study is based on cross-sectional face to face interviews using questionnaires, presented in this paper, consisting of a first part regarding the socio-economic situation and a second part which aimed to classify the disease and any excessive use of drugs.

Of the 610 pharmacists trained with an online course, 446 gathered a total of 4426 correctly compiled questionnaires.

The participation of community pharmacies has highlighted various criticalities especially of an organisational nature; however, it also revealed the power of this method as a means of gathering epidemiological data with a capillarity which few other methods can match.

The objective was also to identify each territory's requirements and facilitate the decision-making process in terms of understanding what patients/citizens actually require.

INTRODUCTION

Migraine headache is a disabling neurovascular syndrome which affects 12-15% of the global population; women are especially prone with a rate of 18-20% among the female population. According to a recently published study in *The Lancet* [1], migraine headache represents the third most commonly cited cause in years lived with disability in both males and females aged 15-49 years.

A review evaluated which other countries in Europe, and in the world, have utilised community pharmacies to identify headache sufferers as a means to make an early identification of the pathology [2].

The early identification of this pathology could potentially lead to enormous savings for the National Health System as well as a marked improvement in the quality of life for the subjects affected by this condition [3].

Among migraineurs, the overuse of medication, a reward seeking behaviour, may be a risk factor in the development of daily migraine and chronic migraine. Indeed, while truly effective medication does exist for the treatment of migraine, its overuse, both in terms of frequency and therapy duration, may have the paradoxical effect of worsening the situation rather than improving it. This condition, called medication overuse headache (MOH), is a growing problem, with an estimated incidence of 1-2% in the general population and up to 30-50% in tertiary headache centres.

Detecting cases of MOH is not straightforward as the majority of migraineurs tend to rely on self-medication with over-the-counter analgesics (OTC) and, in the course of their lifetime, may never turn to a specialist pain centre or even consult their family doctor [2]. Community pharmacists may therefore be in a strategic position to identify and advise individuals who self-medicate, particularly with respect to prevention and early detection of medication overuse to prevent the development of the MOH. Furthermore, they could have an invaluable role by evaluating through simple tests the risk amongst the chronic migraine population of obesity, which leads to significant costs for the healthcare system and induces major disabilities for the affected subject.

Moreover, exploiting the network of community pharmacies makes it possible to identify those "real life" individuals who for socio-economic reasons are unlikely to turn to a tertiary treatment centre such as a headache centre for treatment. This could help to reduce the social divide in terms of those who seek treatment for headaches.

In 2014 our group published the results of a survey conducted in Piedmont, Italy, on the patterns of use and dispensing of drugs in patients requesting assistance from pharmacists for relief of a migraine attack.

The survey was supported by the Italian Headache Foundation (FICEF), in collaboration with the Order of Pharmacists of Turin, Regional Deputy, and the Department of Scienza e Tecnologia del Farmaco, University of Turin.

The study entailed the administration of a 9-item questionnaire, the first part of which comprised items from the ID Migraine Screener test [4, 5] which investigates the different types of headache and whether the subject suffers from migraine headache according to symptom occurrence and severity. On this basis, subjects were categorized as having "Definite migraine" or "Probable migraine".

Thanks to the interesting findings of that study, both from an epidemiological perspective and a public policy viewpoint [6, 7], the decision was taken to expand the scope of the model to a national level in agreement with the Federation of the Orders of Italian Pharmacists (FOFI), the Department of Scienza e Tecnologia del Farmaco, University of Turin (DSTF), the Italian Foundation for Headaches (F.I.C.E.F.), the Epidemiology Unit, ASL TO3, (ASL S.C. a D.J. Servizio Sovrazonale di Epidemiologia), ATF Informatics Society and the Order of Pharmacists of Turin.

MATERIALS AND METHODS

1. DESIGN OF THE STUDY - QUESTIONNAIRE

The study was designed as a *cross-sectional survey* based on face-to-face interviews using questionnaires drawn up by experts and based on the scientific literature; given the unusual nature of the centres responsible for recruiting the subjects, i.e. community pharmacies, it was decided to carry out a specific training course for the data gatherers in order to ensure that the data was collected with the most standardised procedure possible.

2. RECRUITMENT CRITERIA

The subjects were recruited at a maximum rate of 5 patients per pharmacy per month amongst those entering the pharmacy requesting medicine for self-medication of a headache.

3. TRAINING

The training course made use of an on line platform which the pharmacists could log on to by completing a form including their regional code; health service code and AIFA code (see *RECRUITMENT OF PHARMACIES IN THE AREA CREATION OF A PHARMACY DATABASE*). The training consisted of an audio file supported by explanatory slideshow. This standard training course for all territories provided through distance learning methods assured an optimised and uniform training time.

4. RECRUITMENT OF THE PHARMACIES IN THE EXPERIMENTAL AREA - CREATION OF A PHARMACY DATABASE

For this purpose, the AIFA code was used (<http://www.dati.salute.gov.it/dati/dettaglioDataset.jsp?menu=dati&idPac=5> - last date consulted 30/01/2017-). This is an open source system of unique identity codes for each community pharmacy made available by the Ministry of Health.

5. LOGISTICS ORGANISATION

To ensure greater control at the level of the individual provinces, a provincial coordinator was appointed, who reported directly to the head of project coordination.

RESULTS AND DISCUSSION

1. DESIGN OF THE STUDY - QUESTIONNAIRE

With respect to the previous study [6], some modifications to the questionnaire were made (Table 1). In particular, a section regarding the socio-economic situation of the recruited subjects was added (questions 1-6) in order to understand which were the determining factors in the prevalence of headaches.

The socio-economic questions are the product of studies carried out over an extended time span by the research group of the Epidemiology Unit, experienced in the analysis of social disparities in health treatment, [8] and discussions between university, neurologists and pharmacists. These questions have already been included in other studies carried out in community pharmacies with the aim to investigate the correlation between social factors and the pathology under investigation.

Regarding the second part of the questionnaire, which aimed to classify the pathology and any overuse of medication, some new questions were added compared to the previous version used in other studies (Table 2).

No changes were made to the ID Migraine Screener test [4] which investigates the different types of headache and whether the subject suffers from migraine headache according to symptom occurrence and severity. On this basis, subjects were categorized as having "Definite migraine" or "Probable migraine" [5].

The other questions serve to ascertain whether the subject suffering from migraine is in treatment by a medical professional and whether the MOH condition is acute [9]. The overall objective is to optimise the investigative efficiency of this questionnaire.

2. RECRUITMENT CRITERIA

Each pharmacy had the objective to recruit a maximum of 5 subjects/month for each month of the 6-month experiment. Obviously the maximum numerosity of the questionnaire is disproportionate compared to the potential migraineurs, considering the high prevalence of primary headache among the population.

3. TRAINING

Following registration in the database of Pharmacies, the pharmacists underwent a 3-hour training course with a nationally uniform structure for all areas of the country.

The online training course consisted of 3 1-hour sections using a slide presentation:

- 1) classification and epidemiological assessment of primary headaches;
- 2) clinical physio-pathology of primary headaches;
- 3) headache medication and the headache questionnaire.

After completing each section, the pharmacist had to answer 3 multiple-choice questions correctly in order to proceed to the successive section in the training course. Only after passing all 3 sections was the pharmacist deemed to have completed the training course (Table 3-4)

After completing the training procedure, each pharmacist was provided with a unique username and password linked to the Pharmacy database (AIFA code). These could be used to log on to the Questionnaire database in order to access the questionnaire to be filled in online (Tables 1-2). The study was designed in such a way that it was possible to trace the location of the pharmacies and to gather questionnaires systematically so as obtain the maximum amount of location-specific information possible.

4 RECRUITMENT OF PHARMACIES IN THE EXPERIMENTAL AREAS - CREATION OF A PHARMACY DATABASE

Utilising the AIFA code in the Pharmacy database allowed the identification of the location of the pharmacy in Italy. Through the code, it is possible to determine the region, province, city and street address of the pharmacies present in the database.

The system was conceived to reduce the time to a minimum and this allowed rapid access to all the areas involved in the study. In addition to that, the registration process for the pharmacies did not require detailed information which would have made it time-consuming and overly complex. The information requested was, however, sufficient to enable geo-localisation; for the socio-economic stratification, by cross-checking the geographical data and the data gathered in the questionnaires, it was possible to obtain a figure for the evaluation of the requirements.

5 LOGISTICS ORGANISATION

90 provincial coordination officers were appointed thanks to whom it was possible to recruit participants in 59 out of the total of 107 Italian provinces. The number of participating pharmacies was 514, with a total of 610 pharmacists trained (Table 3-B). Of the 610 pharmacists, 446 gathered questionnaires. The total number of correctly compiled questionnaires amounted to 4425. As it is possible to note in Table 4, there is a downward gradient running from north to south of the country both in terms of participating pharmacies/trained pharmacists and the number of questionnaires returned.

Stratification based on socio-economic characteristics is essential in determining how effective the community pharmacy may be in eliminating social disparities

CONCLUSION

The participation of community pharmacies, which was on an entirely voluntary basis, in an area-specific project shed light on various criticalities especially of an organisational nature; however, it also revealed the power of this method as a means of gathering epidemiological data with a capillarity which few other methods can match.

One of the most critical points which must be evidenced is the complexity in creating a pharmacy database: in future, greater use of IT instruments must be made in order to prevent errors in data entry for the pharmacy and personal information.

The use of the AIFA code was an important instrument in tracing the origin of the information entered. However, it became clear that pharmacists had some difficulty in finding their own AIFA code. In any case, this code remains a powerful tool for the future mapping of a territory's requirements based on community pharmacies.

The objective, therefore, was not only to gather data, but to identify each territory's requirements and, hence, facilitate the decision-making process in terms of understanding what patients/citizens actually require.

REFERENCES

1. GBD 2015 Disease and Injury Incidence and Prevalence Collaborators (2016) Global, regional, and national incidence, prevalence, and years lived with disability for 310 diseases and injuries, 1990-2015: a systematic analysis for the Global Burden of Disease Study 2015. *Lancet* 8;385(10053):1545-1602
2. Giaccone M, Baratta F, Allais G, Brusa P (2014) Prevention, education and information: the role of the community pharmacist in the management of headaches *Neurol Sci* 35:S1-4
3. Camarda R, Monastero R (2003) Prevalence of primary headaches in Italian elderly: preliminary data from the Zabùt Aging Project. *Neurol Sci* 24:S122-124

4. Lipton RB, Dodick D, Sadowsky R et al (2003) A self-administered screener for migraine in primary care. The ID Migraine™ validation study. *Neurology* 61:375–382
5. Rapoport AM, Bigal ME (2004) ID-migraine. *Neurol Sci* 25:S258–S260
6. Brusa P, Allais G, Bussone G et al (2014) Migraine attacks in the pharmacy: a survey in Piedmont, Italy *Neuro Sci* 35:S5-9
7. Brusa P, Allais G, Rolando S et al (2015) Migraine attacks in the pharmacy: a gender subanalysis on treatment preferences. *Neurol Sci* 35:S93-95
8. Costa G, Marinacci C, Spadea T (2010) Social inequalities in health: challenges and current priorities in a European context. *Epidemiol Prev* 34(5-6):31-4
9. Saracco MG, Allais G, Tullo V et al (2014) Efficacy of frovatriptan and other triptans in the treatment of acute migraine of normal weight and obese subjects: a review of randomized studies. *Neurol Sci* 35:S115–S119

- 1) **Sex** M/F
- 2) **Age** [actual age]
- 3) **Height** [in cm] and **weight** [in kg]
- 4) **Education** [the highest qualification attained]
 - University degree or higher
 -
 - 4-5-year High School Diploma valid for third-level education
 -
 - 2-3-year High School Diploma not valid for third-level education
 -
 - Junior High School Diploma (or apprenticeship/job)
 -
 - Primary school exam
 -
 - No qualification
- 5) **Occupation**
 - If employed:
 - Manual labourer
 -
 - Employee non-manual (conceptual/intellectual)
 -
 - Self-employed:
 - Businessperson
 -
 - Professional
 -
 - Self-employed worker
 -
 - Unemployed
 -
 - Homemaker
 -
 - Student
 -
 - Retired
 -
 - Other (military service, invalid etc.)
 -
- 6) How many members are there in your family? Number of people in your household (including patient)?

Table 1. "Headache and Community Pharmacies" Questionnaire part 1 -Social Questionnaire.

- 7) **ID Migraine screener test**
 - Did you have the following with your headaches:**
 - You felt nauseated or sick to your stomach? YES/NO
 - Light bothered you (a lot more than when you didn't have headaches)? YES/NO
 - Your headaches limited your ability to work, study, or do what you needed to do for at least 1 day? YES/NO
 - If the patient answers YES to at least three questions, there is a 95% probability that he/she suffers from migraines
- 8) **How long have you been having headaches?** <1 year; 1-4 years; 5-9 years; ≥10 years.
- 9) **How many days have you had a headache in the last 3 months?**n°
- 10) **How many days each month do you take pain medication for a headache?** up to 3; from 4 to 10; more than 10.
- 11) **Does the headache go away after you have taken pain medication?** Yes; Often; Rarely; No.
- 12) **Which medicines do you usually take when you have an acute attack?** NONSTEROIDAL ANTI-INFLAMMATORY DRUGS (NSAIDs); ERGOT-DERIVATIVES; TRYPTANS; COMBINATION DRUGS; OTHER ANALGESICS [more than one answer is permitted].
- 13) **Are you being treated by a medical professional?** [more than one answer is allowed] NOBODY; MY FAMILY DOCTOR; A SPECIALIST; A HEADACHE CENTRE
- 14) **Do you regard your headache as an illness?** YES/NO.

Table 2. "Headache and Community Pharmacy" Questionnaire part 2.